

OPINION OPEN ACCESS

A Pollen-Based Assisted Migration for Rapid Forest Adaptation

David Chludil¹  | Jaroslav Čepl¹  | Arne Steffenrem²  | Jan Stejskal¹  | Christi Sagariya¹  | Torsten Pook³  | Silvio Schueler⁴  | Jiří Korecký¹  | Curt Almqvist⁵  | Debojyoti Chakraborty⁴  | Mats Berlin⁵  | Milan Lstibůrek¹ 

¹Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic | ²Division of Forest and Forest Resources, Norwegian Institute of Bioeconomy Research (NIBIO), Ås, Norway | ³Animal Breeding and Genomics, Wageningen University & Research, Wageningen, The Netherlands | ⁴Department of Forest Growth, Silviculture and Genetics, Austrian Research Centre for Forests (BFW), Vienna, Austria | ⁵Forestry Research Institute of Sweden (Skogforsk), Uppsala Science Park, Uppsala, Sweden

Correspondence: Milan Lstibůrek (lstiburek@fld.czu.cz)

Received: 29 June 2024 | **Revised:** 6 December 2024 | **Accepted:** 12 December 2024

Funding: The research leading to these results has received funding from the EEA/Norway Grants (2014-2021) and the Technology Agency of the Czech Republic.

Keywords: forest nurseries | forestry policy | gene flow | genetic diversity | heterosis | natural selection | phytosanitary risks | seed orchards

ABSTRACT

Climate change poses an unprecedented threat to forest ecosystems, necessitating innovative adaptation strategies. Traditional assisted migration approaches, while promising, face challenges related to environmental constraints, forestry practices, phytosanitary risks, economic barriers, and legal constraints. This has sparked debate within the scientific community, with some advocating for the broader implementation of assisted migration despite these limitations, while others emphasize the importance of local adaptation, which may not keep pace with the rapid rate of climate change. This opinion paper proposes a novel pollen-based assisted migration strategy as a potential middle ground in this debate. By leveraging existing seed orchard infrastructure for controlled pollen transfer, this approach aims to enhance forest resilience through the introduction of genetic material from climatically suitable sources while acknowledging local adaptation. We assess the genetic implications of the proposed strategy through computer simulation. Additionally, we examine the ecological implications of assisted gene flow, discussing the potential benefits of heterosis and the risks of outbreeding depression in intra-specific hybrid populations. We further explore the advantages of pollen-based migration in mitigating phytosanitary risks, reducing economic barriers, and simplifying legal considerations compared to traditional seed or seedling transfer methods. Regional perspectives on adapting pollen-based assisted migration are provided, with specific examples from Northern and Central Europe. We highlight how this approach could be integrated into existing forestry practices and regulatory frameworks within the European Union. We conclude by advocating for the inclusion of pollen-based assisted migration in future international projects and operational forestry, emphasizing the need for adaptable policies that can support innovative forest management strategies in the face of climate change.

1 | Introduction

Forest ecosystems play a crucial role in mitigating climate change through climate regulation (Smith, Baker, and Spracklen 2023) and carbon sequestration, capable of removing billions of tons of carbon from the Earth's atmosphere (Canadell and Raupach 2008).

By fully utilizing available resources, knowledge, and technology, their potential for carbon sequestration could be significantly enhanced (IPCC 2022). However, tree species are particularly vulnerable to the adverse effects of climate change, such as rising temperatures and prolonged droughts, which increase their susceptibility to wildfires, diseases, and pests (Lindner et al. 2010).

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2025 The Author(s). *Global Change Biology* published by John Wiley & Sons Ltd.

These factors contribute to landscape changes and a significant decline in forest cover, impeding forests' natural capacity to absorb carbon. Furthermore, the widespread decline of trees in younger age stages is excessively increasing the amount of released carbon that has been previously stored (Gorte and Sheikh 2010), further elevating atmospheric carbon levels.

Trees can withstand shifts in environmental conditions through phenotypic plasticity (Kramer 1995; West-Eberhard 1989), which enables them to adjust to the inherent diversity of landscapes (Schreiber, Hacke, and Hamann 2015), undergoing continual changes. However, it may fall short of addressing the unforeseeable impacts of climate change (Visser 2008). Therefore, long-term responses such as adaptation through selection and natural migration are essential because an inadequate response to unfavorable conditions could result in species decline (Aitken et al. 2008; Ledig, Rehfeldt, and Jaquish 2012).

Natural or artificial selection is an important process that progressively improves the tolerance to the harmful effects of climate change by increasing the frequencies of beneficial alleles in a population over many generations (White, Adams, and Neale 2007). Tree species have a very long reproductive cycle, which makes this inherently slow adaptation process out of step with the speed of contemporary climate change (Rehfeldt et al. 2002). Thus, migration can be a more relevant climate change mitigating mechanism because it can quickly introduce climatically preadapted genotypes or haplotypes to existing populations (Aitken and Whitlock 2013).

Tree species generally disperse seeds within a limited radius, often just a few tens of meters from the parental tree, thereby gradually increasing their spatial distribution (Burns 1990). However, data shows that forests expanded at an accelerated pace after the last glacial period, which could not have been achieved by typical seed dispersal patterns. This phenomenon, known as Reid's paradox (Reid 1899), challenges our understanding of species expansion. Rare long-distance dispersal events, potentially driven by unusual atmospheric conditions (Clark et al. 1998), may explain the rapid spread.

Pollen-mediated gene flow provides valuable insights into migration dynamics, particularly in wind-pollinated forest tree species. Long-distance pollen dispersal has been repeatedly reported, with well-documented cases of grains traveling hundreds to thousands of kilometers from their source (Campbell et al. 1999; Savolainen, Pyhäjärvi, and Knürr 2007). This emphasizes the critical role of both seed and pollen dispersal in how forest tree species cope with changing environmental conditions, including contemporary climate change. However, models indicate that most tree species' migration rates cannot keep up with the speed of climate change, with some species needing to migrate up to a hundred times faster than historical rates suggest (Sáenz-Romero et al. 2016). On top of that, long-distance dispersion's rate and direction are unpredictable (Nathan et al. 2002, 2011). Contrary to what models predict, the actual migration rates are usually slower, likely due to competition at new sites and low seed germination rates (Sittaro et al. 2017; Zhu, Woodall, and Clark 2012).

Given the limitations of natural migration, there has been a notable focus on human-aided migration, coined by a variety of terms such as assisted migration, assisted colonization (Hoegh-Guldberg et al. 2008), assisted gene flow (Aitken and Whitlock 2013), and managed relocation (Richardson et al. 2009). These conceptual frameworks have been suggested for animals (Bouma et al. 2020) as well as plants (McKone and Hernández 2021), including forest tree species (P. W. Clark et al. 2022). They all share a common goal: to relocate populations to areas with favorable projected climate conditions (Mueller and Hellmann 2008; Ste-Marie et al. 2011).

In this opinion paper, we outline several critical challenges associated with current methods of assisted migration. Importantly, we introduce a novel pollen-based approach to assisted migration, which may offer solutions to these challenges, as outlined in Figure 1. We believe that this innovative method, which could be easily integrated into established frameworks, will significantly contribute to the broader acceptance and effectiveness of assisted migration worldwide.

2 | Current State: General Challenges of Assisted Migration

Assisted migration, the deliberate relocation of tree populations, faces several key challenges (Figure 1) limiting its broader application: (1) environmental factors and current forestry practices, (2) insufficient transfer guidelines, (3) phytosanitary risks from seed and seedling translocation, (4) economic and knowledge barriers faced by local nurseries, and (5) legal obstacles in seed and plant material transport.

Plants' phenotypes evolve to cope with specific local conditions, leading to variations in traits within the same species across their distribution range. A notable meta-analysis revealed that 71% of studies across various organisms reported local adaptations, resulting in an average 45% increase in the fitness of native populations compared to non-native ones (Hereford 2009). Additionally, genomic research supports the idea that local populations are adapted to factors beyond climatic variables, namely soil and photoperiodism (Van Daele, Honnay, and De Kort 2022). Local adaptation is especially prevalent in tree species (Aitken and Bemmels 2016), where differentiated populations are referred to as provenances.

The relative differences in growth and hardiness among provenances have been extensively studied over the past centuries (Langlet 1971). These experiments have primarily aimed to identify the best seed sources for forest establishment by testing the effects of the common environment on different populations. By the late nineteenth century, it was concluded that local provenances usually offer the best balance of growth and hardiness, leading to the preference for local seed sources (Aitken and Bemmels 2016; Langlet 1971). This principle, commonly referred to as "local is best," remains widely accepted. Both traditional forestry focusing on maximizing wood production and close-to-nature practices emphasizing complex natural processes and additional forest functions (O'Hara 2016) continue to prefer the "local is best" principle.




FIGURE 1 | Assisted migration faces numerous challenges, including the unpredictability of local environmental conditions due to climate change, phytosanitary risks associated with the transfer of seeds and live plants, and difficulties faced by stakeholders. Many of these issues can be addressed through a pollen-based approach to assisted migration.

In contrast, earlier recommendations for assisted migration often focused exclusively on climate variables, such as mean annual temperature, while overlooking crucial non-climatic factors like phenological cues and soil properties. These non-climatic factors are highly location-specific and play a vital role in the local adaptation of populations. By ignoring them, one risks overestimating the adaptive potential of assisted migration (Xu et al. 2024; Xu and Prescott 2024). For example, even under optimal climatic conditions, edaphic variables such as soil nutrients, water availability, interactions with soil organisms, and seed predation can significantly limit plant establishment in new areas (Benning and Moeller 2021; Brown and Vellend 2014; Ni and Vellend 2024). Phenological cues are equally important to the success of assisted migration, especially when relocating tree populations over long distances. Many species rely on site-specific day length, rather than temperature, to regulate phenological events (Basler and Körner 2012; Laube et al. 2014; Way and Montgomery 2015). Consequently, translocating these species to areas with different day lengths can cause mismatches in the timing of phenological events, leading to reduced growth and increased vulnerability to spring and autumn frosts (Montwé et al. 2018; Silvestro et al. 2019). This reduces the distances over which populations can be relocated, making it more challenging to align assisted migration efforts with the pace of climate change (Cooper et al. 2019; Stinziano and Way 2014).

However, recent advancements in the collection and availability of high-quality, large-scale landscape data (Dauphin et al. 2023) are addressing these limitations by incorporating a broader range of environmental variables into assisted migration frameworks (Feng et al. 2020). This holistic approach not only aligns with traditional forestry practices that emphasize local adaptation but also offers a more robust basis for assisted migration. Continued research in this field is essential to develop robust recommendations that account for both climatic and non-climatic variables, thereby reducing the risk of maladaptation and promoting population fitness in new environments.

Another significant risk of assisted migration is the potential introduction of non-native pests and pathogens. This risk increases with the growing volume of international trade in plants, including trees (Eschen, Roques, and Santini 2015; Liebhold et al. 2012). As a result, only a small fraction of imported plants undergo inspection (Brasier 2008). Furthermore, the effectiveness of these inspections is limited because infested plants may not show symptoms upon arrival, making it difficult to ensure complete phytosanitary safety through traditional methods (Jung et al. 2016; Liebhold et al. 2012). Phytosanitary risks are not confined to living plants alone. Recent studies suggest that seeds also pose a significant phytosanitary risk, despite previously being considered relatively safe for international transfer. This risk is particularly pronounced in Europe, where regulations for preventing the introduction of seed-borne pathogens are less stringent compared to other regions (Franić et al. 2019; Vettraino, Potting, and Raposo 2018).

Forest nurseries are crucial for forest regeneration, requiring increased seedling production and diversity to address contemporary climate change challenges in forestry. A North American study (P. W. Clark et al. 2023) highlights the need for more ecological diversity in nurseries, as the focus on commercially valuable species often limits conservation and climate adaptation efforts. While there are business opportunities to produce seedlings adapted to future climatic conditions, including those phenotypes resistant to climate extremes and new pests, this bears additional costs, logistical complexities, and financial risks. Nurseries often prioritize uniformity in seed lot germination and growth for easier cultivation and more effective production (Ivetić et al. 2016), which often contrasts with the desired level of genetic diversity.

These challenges, especially when sourcing seeds from distant locations, have hampered investments, particularly in regions where the number of nurseries has declined in recent decades in Europe and North America. Moreover, small nurseries fear

competition from larger foreign entities threatening their profitability in an open-market scenario. Nursery managers across Central Europe have also expressed concerns about the impacts of climate change on their operations. Perceptions vary by country, with Austria and Germany expecting higher demand for non-native species, while Poland and Hungary anticipate a need for more resilient local varieties (Hazarika et al. 2021).

Finally, implementing assisted migration requires considering various legal regulations, including the “Nagoya Protocol” (Secretariat of the Convention on Biological Diversity 2011). It may constrain assisted migration in countries that have adopted this international agreement with stringency. Nevertheless, assisted migration is predominantly hindered by laws on a national and regional level, restricting seed usage to specific “seed zones” or “provenance regions”, thereby favoring locally adapted populations (Konnert et al. 2015). Thus, a dynamic policy framework adapting to diverse climate change scenarios is essential for ensuring long-term forest health and resilience.

3 | Assisted Gene Flow: Implications of Inter-Provenance Hybridization

Assisted migration methodologies can be categorized into two general approaches: migration outside the species' geographic range and migration within it. It is anticipated that assisted migration of tree species will predominantly occur within the species' geographic range (Sáenz-Romero et al. 2021). This approach poses less ecological risk as it avoids relocating trees to unfamiliar environments where unforeseen interactions could occur (Aitken and Whitlock 2013). Additionally, it is generally more accepted by the public compared to the alternative (Findlater et al. 2020).

Within-range assisted migration has been proposed by Aitken and Whitlock (2013) and Kelly and Phillips (2016). These studies suggest leveraging existing genetic differences among tree populations of a given species by transferring seeds or seedlings to introduce preadapted genotypes and increase adaptive genetic variation of endangered populations, through “assisted gene flow.” However, as these authors caution, this strategy carries the risk of unintended interspecific hybridization, particularly with closely related species, leading to genetic incompatibilities (Critchfield 1967). Furthermore, the effectiveness of genetic infusion of preadapted genotypes can be limited by potential mismatches in flowering phenology and will only manifest once the introduced seedlings reach reproductive age.

The timing of flowering events in plants is generally highly variable (Shelton et al. 2024; Weis 2015). While this variability is more pronounced in herbaceous species, it is also observed in tree species, albeit to a lesser extent (Horbach, Rauschkolb, and Römermann 2023). This variation is a result of adaptation to local conditions, which can occur in a relatively few generations. Differences in flowering phenology may result in flowering asynchrony both within and among populations. This asynchrony potentially increases the current variation in the reproductive success of forest tree species, thus affecting genetic diversity. Flowering phenology asynchrony tends to increase with geographic distance, possibly resulting in phenological

isolation of populations. As a result, migrants rarely mate with members of the resident population, promoting assortative mating patterns, with a higher rate of mating occurring between migrants and between residents (Wadgymar and Weis 2017). Even when mating between migrants and residents is successful, cones or fruits may not be produced by late-flowering individuals (Wadgymar, Cumming, and Weis 2015). This leads to a reduced rate of hybridization that becomes even more apparent in the second generation of hybrids (F_2) (Wadgymar and Weis 2017).

Apart from flowering asynchrony, assisted gene flow may lead to emergent short-term genetic effects resulting from the hybridization of genetically divergent populations, which might hinder achieving the desired adaptation. These effects can include potential positive impacts such as heterosis in the first generation of hybrids or negative consequences like outbreeding depression and hybrid decay in subsequent generations.

Heterosis, also frequently called hybrid vigor, is the beneficial effect that occurs when a hybrid surpasses the quality of both parents (Shull 1914). This phenomenon was already recognized by Charles Darwin (1876) and is traditionally exploited in agriculture to enhance the yield of inbred crop lines and improve resistance to diseases (e.g., Hei, Hussein, and Laing 2016). The degree of heterosis generally rises with increasing genetic distance up to a certain point, after which it declines (e.g., Moll et al. 1965; Würschum et al. 2023). Heterosis is also apparent in inter-species crosses and is utilized in forestry species, such as in the larch genus, with European larch (*Larix decidua* Mill.) and Japanese larch (*Larix kaempferi* (Lamb.) Carr.). First-generation (F_1) larch hybrid (*Larix* × *eurolepis* Henry) has shown significant genetic response in growth rate, wood quality, and seed yield (Marchal et al. 2017). Thus, they pose overall better performance compared to the parent species and also compared to the second-generation hybrids (F_2). Similar results have been found in poplar (Zanewich, Pearce, and Rood 2018), eucalyptus (Madhibha et al. 2013), silver fir (Stejskal, Horák, and Typta 2016), and pines (Dungey 2001).

There are two prevalent hypotheses that explain the basis of heterosis. The first is the dominance hypothesis, which suggests that in hybrids, harmful recessive alleles are hidden by beneficial alleles from either parent (Bruce 1910; Davenport 1908; Jones 1917). The second is the overdominance hypothesis, which proposes that heterozygosity itself can result in hybrids being more vigorous than either parent (East 1908; Shull 1908). Those approaches were further expanded by the epistatic theory of heterosis, involving interactions among genes (Powers 1944). However, the exact genetic and molecular basis behind heterosis is still not fully understood (Birchler et al. 2010; Yu et al. 2021).

The question arises whether heterosis occurs only in isolated groups, such as between inbred lines and within inter-specific hybrids, or also across populations connected by gene flow, similarly to forest trees. Simulations of these scenarios suggest that heterosis can indeed occur in such cases (Whitlock et al. 2000). Moreover, various forest tree species have been tested for potential heterosis effects in their provenance hybrids (Liesebach, Liepe, and Bäucker 2021). Harfouche and Kremer (2000), Harfouche et al. (2000) reported an 8.4% increase in growth due

to heterosis in maritime pine (*Pinus pinaster* Aiton) provenance hybrids. Additionally, Doerksen, Bousquet, and Beaulieu (2014) revealed a heterosis effect in inter-provenance crosses of white spruce (*Picea glauca* (Moench) Voss), with approximately 6% greater height growth at age 15 compared to intra-provenance progenies. In contrast, Eriksson and Ilstedt (1986) observed heterosis in Norway spruce (*Picea abies* (L.) H. Karst) provenance hybrids at only one of three test sites, while Kaya and Lindgren (1992) found no evidence of heterosis. Similarly, Levkoev et al. (2017) did not observe an increase in growth and wood density in hybrids compared to the local population. Although heterosis was observed in several intra-specific scenarios, its impact does not seem to be as pronounced as in the inter-species hybrids of forest trees.

Conversely, outbreeding depression occurs in hybrids of distant populations when local adaptive traits average out, resulting in lower fitness in later generations compared to the parental lineage. Although common in plants (Liesebach, Liepe, and Bäucker 2021), it is rare in forest trees, with Sakhalin fir (*Abies sachalinensis* F. Schmidt) being a single case study (Goto et al. 2011). Outbreeding depression typically appears from the second generation onwards, as hybrid vigor masks these effects in the first generation. From the F_2 generation, recombination disrupts epistatic interactions between coadapted genes (Whitlock et al. 2013). However, these interactions are expected to be restored over several generations, and the fitness of later generations can surpass that of the parental generation as climatically preadapted alleles will become apparent (Aitken and Whitlock 2013; Grummer et al. 2022).

Understanding the genetic architecture of adaptive traits is crucial for evaluating the benefits and risks of assisted gene flow such as the aforementioned heterosis and outbreeding depression. This involves examining the number, position, and effect sizes of individual genes. Genome-wide association studies provide detailed molecular-level insights into gene-trait associations. While detecting gene effects is relatively efficient in inbred lines, it becomes significantly more complex in outbred populations (Lynch and Walsh 1998). Outbred forest tree species present unique difficulties due to their large genomes and growth in highly heterogeneous environments.

Empirical evidence from tree species indicates that adaptive traits, such as resistance to drought, frost damage, and pathogens, often exhibit polygenic or complex architectures (De La Torre et al. 2019, 2021; Depardieu et al. 2021; Milesi et al. 2019; Riehl et al. 2023; Singh et al. 2024). These architectures may consist of a few genes with relatively large effects but are largely controlled by many genes with smaller effects. This predominantly polygenic inheritance of quantitative traits in forest trees is further supported by traditional genetic variance decomposition, based on controlled crosses and progeny trial evaluations (White, Adams, and Neale 2007).

4 | Pollen-Based Assisted Migration as an Innovative Approach

We propose a comprehensive strategy framed around intentional pollen transfer from suitable sources (typically referred

to as a population of migrants) to reduce the risk of maladaptation in adverse climates while preserving the integrity of locally adapted populations. This strategy involves collecting pollen from a source seed orchard and transferring it to a target (typically referred to as a resident population) seed orchard for intraspecific hybridization (Figure 2), thereby introducing haplotypes suited for anticipated future climate conditions.

A key advantage of this method is that it capitalizes on existing forestry infrastructure, specifically, seed orchards, which already play a critical role in global forestry for large-scale seed production (White, Adams, and Neale 2007). In Europe specifically, the establishment of seed orchards for seed mass production began in the 1950s (see Faulkner (1975), for early review) and has continued to grow, with the current number exceeding 1500 seed orchards of 40 forest tree species (EU Forest Reproductive Material Information System, <https://ec.europa.eu/forematis>). Traditionally, seed orchards have been closely linked to tree improvement programs, serving as the primary source for producing bulk seed for afforestation and reforestation efforts. Many seed orchards are also established for tree species of lower economic but significant ecological importance (e.g., So, Theilade, and Dell 2010). Globally, seed orchard networks extend beyond Europe to North America (Li, McKeand, and Weir 1999; Miller and DeBell 2013; Reid 2008), South America (Jayawickrama and Balocchi 1993; Rockwood, Huber, and White 2001), Africa (Gapare 2000; Missanjo, Kamanga-Thole, and Manda 2013; Swain, Verryin, and Laing 2013), Asia (Batkhue et al. 2010; Chaix et al. 2011; Leksono, Kurinobu, and Ide 2008; Moriguchi et al. 2005; Na et al. 2015; Nicodemus et al. 2009; Shi et al. 2016), Australia (Burczyk et al. 2002; Johnson, Robinson, and O'Hara 1992; Wu et al. 2007), and New Zealand (Burdon, Carson, and Shelbourne 2008; Dickson, Sweet, and Mitchell 2000; Turner 1997).

Collecting pollen in the source orchard is feasible because the trees in orchards are pruned to facilitate seed harvest and control pollination. When choosing a deployment strategy, several suitable source seed orchards can be identified to obtain the desired pollen mixture. Thus, the pollen mixture could be adjusted frequently based on previous experience and the progression of climate change.

Since open pollination (natural random mating) occurs in seed orchards, implementing supplemental mass pollination (SMP) or controlled mass pollination (CMP) in the target orchard is straightforward. SMP or CMP involves disseminating the collected pollen during the peak receptivity of female strobili (Funda and el-Kassaby 2012), promoting gametic exchange between migrant and resident population genotypes. Additionally, SMP or CMP could be combined with the method of delayed blooming, which adjusts flowering times to promote synchronization. This reduces pollen contamination from inferior genotypes in non-selected populations near the seed orchard (Korecký and El-Kassaby 2016; Song et al. 2018) and increases the annual seed set (Pearse et al. 2015), resulting in higher seed orchard production.

In the proposed strategy, seeds produced in the target seed orchard will be sown in local nurseries (Figure 2), allowing progenies to undergo their entire embryogenesis and early growth under local conditions. This contrasts with traditional transfer




FIGURE 2 | The map is derived from the species distribution model developed by Chakraborty et al. (2021). The green gradient represents the probability of Norway spruce distribution in Europe within the 2041–2060 time range under intermediate climate scenario RCP 4.5. Brown dots show the locations of individual seed orchards across Europe based on the European Union's FOREMATIS database and the internal Norwegian database of seed orchards. The flowchart represents the transfer of desired haplotypes from source to target seed orchards through pollen and subsequent seedling production utilized in target reforestation sites. Map lines delineate study areas and do not necessarily depict accepted national boundaries.

methods where seeds or mature plants carry their original epigenetic profiles when moved to new locations.

Epigenetics, transgenerational changes that modulate phenotype without altering the genetic composition, is increasingly recognized as an important factor in shaping adaptive traits in plant breeding and ecosystem conservation (Bräutigam et al. 2013). Although the effects of epigenetics on tree species remain understudied, its significant role has been suggested in both Norway spruce (Johnsen et al. 2005, 2009) and Scots pine (Bose et al. 2020). Our method may present a significant advantage, as local conditions can modulate gene expression through various epigenetic mechanisms (Franks and Hoffmann 2012), thus enhancing the adaptive potential of these tree species in their native environments and reducing the risk of transferring potentially harmful epigenetic settings.

The pollen-based approach facilitates the production of hybrid offspring with flowering periods that average the flowering times of their parents. Empirical evidence supports this pattern, particularly in fruit trees like apples (Urrestarazu et al. 2017) and *Prunus* species, such as sweet cherries and almonds (Branchereau et al. 2022, 2023; Castède et al. 2014; Dirlewanger et al. 2012; Sánchez-Pérez et al. 2014). Similar trends have also been observed in outcrossing plants like maize (Buckler et al. 2009). These studies reveal that flowering periods are highly heritable and governed by complex genetic architecture. Moreover, Urrestarazu et al. (2017) highlight that the genes controlling flowering phenotype predominantly exhibit additive effects.

The flowering time of hybrids that more closely align with the resident population will gradually increase the proportion of backcrosses and later-generation hybrids within the population.

This, in turn, will boost the frequencies of pre-adapted alleles in subsequent generations. This process can be particularly advantageous for enhancing adaptive genetic diversity, as it establishes a solid foundation for natural selection by favoring the best-adapted individuals among a broad range of genotypes, especially under the competitive pressures of a changing climate. To achieve this objective, we recommend supplemental planting of orchard-derived hybrid seedlings in a grid or group planting (Saha et al. 2012) among the naturally grown seedlings in situ.

This approach aligns with close-to-nature forestry principles, where forest stands are primarily restocked by trees growing from seeds that fall and germinate on-site, maximizing the natural processes within the stand (O'Hara 2016). However, in cases where significant heterosis is present, hybrids can surpass all local individuals in fitness, reducing the proportion of naturally regenerated individuals in favor of F_1 hybrids. This poses a risk, as the initial heterosis diminishes in the F_2 generation, leading to a significant decline in overall fitness and resulting in detrimental effects on the stand (Kaya and Lindgren 1992). In such scenarios, a more effective strategy might be to fully utilize the benefits of heterosis in F_1 hybrids by establishing pure hybrid stands and regenerating them through clear cuts. The establishment of pure hybrid stands can lead to increased biomass production and overall forest productivity due to the superior growth characteristics and enhanced resistance to environmental stresses of F_1 hybrids. This method ensures that the hybrids' genetic advantages are fully exploited during their peak growth periods, maximizing yield and economic return. Additionally, this strategy still facilitates the introgression of climatically pre-adapted alleles, as their pollen contribution to the neighboring stands would promote backcrossing, gradually introducing pre-adapted

alleles and thereby maintaining the overall fitness. Since this process is more gradual, it helps to mitigate the negative effects of hybrid decay in F_2 .

The pollen-based method can also facilitate the adaptation of exotic species. Douglas fir (*Pseudotsuga menziesii* (Mirbel) Franco), native to the Pacific Northwest of North America, is a prime example of an introduced species in Europe. Breeding programs for Douglas fir are already established in several European countries, with seed orchards located in Austria, France, Germany, and other regions. In some countries, such as the Czech Republic and Slovenia, these programs are just being established. Another example of a successful introduction is a Turkish fir (*Abies bornmülleriana* Mattf.) and Nordmann fir (*Abies nordmanniana* (Steven) Spach), with seed orchards in Denmark (Nielsen, Xu, and Hansen 2020; Xu, Nielsen, and Hansen 2018).

These exotic species typically underwent intense selection for both adaptation and productivity traits. While the genetic diversity of mature stands after introduction seems to be comparable with the native populations, it can be reduced due to population bottlenecks caused by the variation in reproductive success, which is a specific problem in exotic species as their populations in non-native environments are typically fragmented (Neophytou, van Loo, and Hasenauer 2020). The pollen-based strategy enhances local populations by increasing genetic diversity and introducing climate-adaptive alleles from suitable regions within the native range.

Beyond the aforementioned benefits for native and exotic species, the pollen-based assisted migration strategy offers solutions to remaining challenges, including phytosanitary risks, economic obstacles, and legislative constraints. Managing phytosanitary risks becomes simpler as pollen is typically unsuceptible to pathogens and is expected to be germinated on site. The approach is economically feasible as seed orchard managers could handle a local pollen collection and storage framework. Nevertheless, some investment in pollen storage may be necessary. As both seed and seedlings are locally sourced, forest nurseries can continue business operations without disruption. Considering these advantages, we anticipate a wider acceptance of the pollen-based assisted migration, leading to legislative adaptations enabling pollen transport.

5 | Simulation Insights Into Genetics of an Adaptation Through Pollen-Based Assisted Migration

From a genetic perspective, the effectiveness of the proposed pollen-based assisted migration depends on several factors. These include the degree of population divergence, as reflected by differences in genetic structure (allelic and genotypic frequencies) and fitness distribution between the source and target populations under current environmental conditions; the extent of gametic phase disequilibrium; the genetic architecture of fitness traits (including the number of loci contributing to trait variance and their respective additive, dominance, and epistatic effects); biological mechanisms of heterosis and

outbreeding depression; narrow-sense heritability; and the intensity of pollen migration from the source orchard to the target orchard.

To examine the genetic implications of pollen-based assisted migration, we utilized a computer simulation model presented in Supplement S1, which includes parametrization and computer code. A schematic overview of the simulation scenarios is provided in Figure 3. Comparative results for pollen-based assisted migration, conventional assisted migration, and a reference scenario relying solely on local populations are shown in Figure 4. The model incorporates a mixed genetic architecture, with 30% of the genetic variance attributed to 10 quantitative trait loci (QTLs), while the remaining genetic variance is polygenic. Narrow-sense heritability was set at 0.2, reflecting a significant environmental component in adaptive traits. The model is specifically designed to simulate diploid populations and, in its current implementation, does not support the evaluation of interspecific hybridization.

We first evaluated a single reproductive outcome in the target seed orchard subjected to pollen-assisted migration, with fitness assessed in the offspring population (forest stands originating from the orchard). Our model revealed significant initial heterosis, resulting in a mean fitness that exceeded the average of parental populations, and even surpassed the mean fitness of conventional assisted migration (generation 1 in Figure 4). However, we should note that the initial population divergence (difference between the allelic frequency of the two founder populations) was assumed to be very large which promoted the initial heterosis and outbreeding depression. Typically, differences between allelic frequencies would be lower, thus the mean fitness would more closely resemble the average of parental populations in the same figure. However, the real adaptive response of the pollen-based assisted migration might be underestimated in our simulations given additional components, such as retaining the local adaptation when using the seed orchard in the target location and additional beneficial factors that were discussed in Section 4.

The dynamic nature of climate change necessitates frequent recalibration of assisted migration models and optimization of pollen mixtures. This also highlights the importance of periodical updates of the genetic composition of seed orchards in both source and target populations adopting a dynamic system that leverages both local (e.g., Lstibůrek, García-Gil, and Steffenrem 2023) and distant adaptation, ensuring proper synchronization with practical forest management as discussed in the previous chapter.

In line with these options, Figure 4 also illustrates the potential for maintaining forest plantations established in generation 1 and allowing natural regeneration in subsequent generations. In generation 2, the model predicts an initial outbreeding depression from reduced initial heterosis and unfavorable epistatic effects. In subsequent generations, however, outbreeding depression is purged and the higher overall genetic diversity in the population leads to higher subsequent adaptive response, eventually resulting in mean fitness comparable to that of conventional assisted migration (assuming the population remains in situ and continues to reproduce locally).




FIGURE 3 | Schematic representation of the computer simulation process. The simulation begins with the creation of two founder populations, distinguished by allelic frequencies: One in the source location (light red background) and the other in the target location (light blue background). Thirty generations of random mating with a constant population size of 5000 individuals facilitated the build up of linkage disequilibrium (LD). The source founder population was modeled with a higher mean fitness compared to the target population, indicating adaptive superiority when introduced to the target environmental conditions. Subsequently, a seed orchard containing 100 individuals was established in each location, derived from their respective founder populations. Assisted migration was then simulated, wherein random mating occurred in the source seed orchard, and a forest stand of 5000 trees was established in the target location (generation 1). From this stand, 500 trees with superior phenotypes (top fitness) were selected and subsequently randomly mated to generate 5000 offspring. Further generations of natural regeneration (2–5) maintained a population size of 5000 individuals through successive cycles of natural selection. This process is represented by the leftmost forest stands in the target location (darker red ellipses). Alternatively, an additional forest stand was established via assisted pollen migration. In this scenario, male haplotypes from the source seed orchard were combined with female haplotypes from the target seed orchard. The resulting forest stand underwent similar mating and natural selection processes up to generation 5, depicted by the middle forest stands in the target location (darker red/blue split-color ellipses to signify mixed haplotypes). For reference, a control scenario without assisted migration was simulated, where random mating occurred within the target seed orchard. This scenario is represented by the rightmost forest stands in the target location (darker blue ellipses).

While we aimed to align simulation parameters with empirical evidence, caution is warranted in generalizing our findings across species. We propose systematic computer modelling research to explore complex dynamics of the proposed system. Fine-tuning simulation parameters for specific species is necessary to evaluate feasibility and account for additional consideration, including benefits of pollen-based assisted migration discussed in our manuscript but not incorporated into the current simulation.

6 | Regional Perspective: Adapting Pollen-Based Assisted Migration in Northern and Central Europe

The Scandinavian region has a long history of successfully introducing and transferring seed sources of Norway spruce and Scots pine (*Pinus sylvestris* L.) (Myking et al. 2016).

Consequently, both selectively bred populations and commercially produced seed lots contain a significant proportion of non-local genotypes transferred from various regions within and between Northern European countries and beyond. Given this established practice, current official transfer guidelines are based on past provenance testing and practical regeneration experience, often tolerating substantial transfers from local origin.

New transfer effect models have been developed for both species to predict performance in future climates and facilitate assisted migration (Berlin et al. 2016; Liziniewicz et al. 2023). These models utilize tree phenotype data from all Scandinavian countries to provide uniform deployment recommendations across the region and to develop joint decision support tools. Implementing such a framework requires addressing national transfer rules, legal barriers, and differences in seed orchard definitions.




FIGURE 4 | Predicted mean fitness values in the target location based on computer simulation. The three scenarios include: (1) seedling-based assisted migration (red circles), where seedlings derived from the source seed orchard are planted in the target location; (2) pollen-based assisted migration (violet squares), where pollen is transferred from the source orchard to the target orchard, and the resulting offspring are planted in the target location; and (3) local offspring (blue circles), where offspring derived from the orchard in the target location are planted in the same area. Generation 1 represents forest stands established using seeds collected from the target seed orchard for each respective scenario. Generations 2–5 represent hypothesized forest stands resulting from natural mating among the top 500 individuals in each stand.

For Scots pine, common deployment recommendations have been successfully developed and officially approved in Sweden and Finland (Berlin et al. 2019). Additionally, a joint Swedish-Finnish decision support tool called “Plantval” (Skogforsk 2024) aids in selecting optimal seed sources. The tool considers the origin of both the seed orchard parents and the pollen, which is assumed to come from the orchard itself or surrounding populations (e.g., Heuchel et al. 2022). Introducing external pollen with known origin and genetic gain through a pollen-based approach can easily be integrated into the tool, directly influencing the model’s predictions and deployment recommendations.

In the Nordic region, seed orchards have undergone genetic thinning and selective harvesting to enhance adaptive properties, increase tolerance to pests/diseases (e.g., Scots pine blister rust and root rot), and boost genetic gain. However, these

methods are limited to existing orchard genotypes. A pollen-based approach would enable a rapid and substantial improvement in one or more desired traits, particularly if those traits are not achievable with the current genetic variation within the orchard.

In contrast to Northern Europe, Central Europe encompasses a broader range of ecoregions and forest types, from xerophytic mixed oak forests to subalpine mountain forests, providing habitat for a diverse array of woody plant species. Forest genetic resource management in Central European countries has traditionally focused on selecting and transferring genetic material with superior growth characteristics, as well as the conservation of major and rare tree species. Notably, conifers such as European Larch and Norway spruce have been historically transferred over larger distances, often outside their native range

(Jansen, Konrad, and Geburek 2017; Jansen and Geburek 2016). However, rare and scattered tree species were often neglected in historical forest management practices, potentially impacting their genetic variation (Hemery et al. 2010; Lefèvre 2004).

Genetic studies on many Central European species have revealed that populations south of the Alps, in refugial areas, exhibit the highest genetic differentiation, while populations north of the Alps display the highest genetic diversity, likely due to the admixture of lineages during recolonization (Petit et al. 2003). Awareness of genetic resources, particularly for scattered tree species, increased towards the end of the 20th century, leading to the establishment of several national gene conservation programs (Müller 1997; Tabel 1997) and the EUFORGEN Network for scattered broadleaves. Additionally, the EC directive for forest reproductive material (1999/105/EC) was enacted, regulating the management of 47 tree species across Europe. These efforts have resulted in the establishment of numerous seed orchards for gene conservation.

Given anticipated shifts in tree species composition due to climate change, there is an increasing need to diversify tree species mixtures and incorporate drought-tolerant species from continental or Mediterranean climates. Central Europe has over a century of experience in assisted migration. One example could be a successful transfer of exceptional “Slavonian oaks”, valued for their high genetic quality and adaptability, from the Save and Drava river plains in Southeastern Europe to Northwestern Germany (Gailing et al. 2012). Similarly, “Calabrian” silver firs from South Italy have been successfully used for reforestation in Denmark (Larsen 1981). Extensive provenance experiments for various tree species were conducted in Europe during the 20th century, and recent comprehensive analyses of these experiments have led to transnational transfer guidelines (Chakraborty et al. 2024). Central European countries can leverage their existing network of 717 seed orchards (encompassing 23 native species and hybrids) to implement pollen-based assisted migration. These orchards are expected to serve as both pollen sources and recipients across a wide range of climatic conditions.

7 | Conclusions

We propose a pollen-based assisted migration strategy as an effective and scalable solution for adapting forests to climate change. Through reasoned examples, we discuss the potential benefits and practicalities of implementing this approach in Scandinavia and Central Europe and its compatibility with broader EU frameworks. We strongly advocate for its consideration in future international projects and operational forestry. A dynamic policy framework adaptable to diverse climate scenarios is essential to fully capitalize on this innovative strategy. Such a framework could serve as a catalyst for revising current EU regulations, thereby enabling more agile and effective forest adaptation strategies, crucial for long-term societal resilience against climate change. Since pollen-based assisted migration relies methodically on seed orchards, we advocate for conventional assisted migration in species and regions where seed orchards are unavailable.

Author Contributions

David Chludil: conceptualization, data curation, investigation, visualization, writing – original draft. **Jaroslav Čepl:** formal analysis, software, visualization, writing – review and editing. **Arne Steffenrem:** investigation, resources, writing – review and editing. **Jan Stejskal:** data curation, writing – review and editing. **Christi Sagariya:** data curation, formal analysis, software. **Torsten Pook:** formal analysis, resources, software. **Silvio Schueler:** writing – review and editing. **Jiří Korecký:** writing – review and editing. **Curt Almqvist:** writing – review and editing. **Debojyoti Chakraborty:** writing – review and editing. **Mats Berlin:** writing – review and editing. **Milan Lstibůrek:** conceptualization, formal analysis, investigation, supervision, writing – review and editing.

Acknowledgments

We thank the anonymous reviewer for valuable suggestions, which have significantly enhanced the quality of the manuscript. The research leading to these results has received funding from the EEA/Norway Grants 2014–2021 and the Technology Agency of the Czech Republic.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data and code that support the findings of this study are openly available in Figshare at <https://doi.org/10.6084/m9.figshare.28033913>. Due to privacy concerns, the locations of the Norwegian seed orchards are not publicly available, but can be obtained by contacting Dr. Arne Steffenrem at arne.steffenrem@nibio.no. The species distribution model can be accessed from Zenodo at <http://doi.org/10.5281/zenodo.3686918>.

References

- Aitken, S. N., and J. B. Bemmels. 2016. “Time to Get Moving: Assisted Gene Flow of Forest Trees.” *Evolutionary Applications* 9, no. 1: 271–290. <https://doi.org/10.1111/eva.12293>.
- Aitken, S. N., and M. C. Whitlock. 2013. “Assisted Gene Flow to Facilitate Local Adaptation to Climate Change.” *Annual Review of Ecology, Evolution, and Systematics* 44: 367–388. <https://doi.org/10.1146/annurev-ecolsys-110512-135747>.
- Aitken, S. N., S. Yeaman, J. A. Holliday, T. Wang, and S. Curtis-McLane. 2008. “Adaptation, Migration or Extirpation: Climate Change Outcomes for Tree Populations.” *Evolutionary Applications* 1, no. 1: 95–111. <https://doi.org/10.1111/j.1752-4571.2007.00013.x>.
- Basler, D., and C. Körner. 2012. “Photoperiod Sensitivity of Bud Burst in 14 Temperate Forest Tree Species.” *Agricultural and Forest Meteorology* 165: 73–81. <https://doi.org/10.1016/j.agrformet.2012.06.001>.
- Batkhuu, N.-O., D. K. Lee, J. Tsogtbaatar, and Y. D. Park. 2010. “Seed Quality of Siberian Larch (*Larix sibirica* Ldb.) From Geographically Diverse Seed Sources in Mongolia.” *Scandinavian Journal of Forest Research* 25, no. sup8: 101–108. <https://doi.org/10.1080/02827581.2010.485815>.
- Benning, J. W., and D. A. Moeller. 2021. “Plant–Soil Interactions Limit Lifetime Fitness Outside a Native Plant’s Geographic Range Margin.” *Ecology* 102, no. 3: e03254. <https://doi.org/10.1002/ecy.3254>.
- Berlin, M., C. Almqvist, M. Haapanen, et al. 2019. *Common Scots Pine Deployment Recommendations for Sweden and Finland*. Uppsala: Skogforsk Report. https://www.skogforsk.se/cd_20190510095857/contentassets/d6472db34bec4f10b0c9b5c694f2fd08/arbetsrapport-1017-2019.pdf.

- Berlin, M., T. Persson, G. Jansson, et al. 2016. "Scots Pine Transfer Effect Models for Growth and Survival in Sweden and Finland." *Silva Fennica* 50, no. 3: 1562. <https://doi.org/10.14214/sf.1562>.
- Birchler, J. A., H. Yao, S. Chudalayandi, D. Vaiman, and R. A. Veitia. 2010. "Heterosis." *Plant Cell* 22, no. 7: 2105–2112. <https://doi.org/10.1105/tpc.110.076133>.
- Bose, A. K., B. Moser, A. Rigling, et al. 2020. "Memory of Environmental Conditions Across Generations Affects the Acclimation Potential of Scots Pine." *Plant, Cell & Environment* 43, no. 5: 1288–1299. <https://doi.org/10.1111/pce.13729>.
- Bouma, A., G. Kuchling, S. Y. Zhai, and N. Mitchell. 2020. "Assisted Colonisation Trials for the Western Swamp Turtle Show That Juveniles Can Grow in Cooler and Wetter Climates." *Endangered Species Research* 43: 75–88. <https://doi.org/10.3354/ESR01053>.
- Branchereau, C., C. Hardner, E. Dirlewanger, et al. 2023. "Genotype-By-Environment and QTL-By-Environment Interactions in Sweet Cherry (*Prunus avium* L.) for Flowering Date." *Frontiers in Plant Science* 14: 1142974. <https://doi.org/10.3389/fpls.2023.1142974>.
- Branchereau, C., J. Quero-García, N. H. Zaracho-Echagüe, et al. 2022. "New Insights Into Flowering Date in *Prunus*: Fine Mapping of a Major QTL in Sweet Cherry." *Horticulture Research* 9: uhac042. <https://doi.org/10.1093/hr/uhac042>.
- Brasier, C. M. 2008. "The Biosecurity Threat to the UK and Global Environment From International Trade in Plants." *Plant Pathology* 57, no. 5: 792–808. <https://doi.org/10.1111/j.1365-3059.2008.01886.x>.
- Bräutigam, K., K. J. Vining, C. Lafon-Placette, et al. 2013. "Epigenetic Regulation of Adaptive Responses of Forest Tree Species to the Environment." *Ecology and Evolution* 3, no. 2: 399–415. <https://doi.org/10.1002/ece3.461>.
- Brown, C. D., and M. Vellend. 2014. "Non-Climatic Constraints on Upper Elevational Plant Range Expansion Under Climate Change." *Proceedings of the Royal Society B: Biological Sciences* 281, no. 1794: 20141779. <https://doi.org/10.1098/rspb.2014.1779>.
- Bruce, A. B. 1910. "The Mendelian Theory of Heredity and the Augmentation of Vigor." *Science* 32, no. 827: 627–628. <https://doi.org/10.1126/science.32.827.627.b>.
- Buckler, E. S., J. B. Holland, P. J. Bradbury, et al. 2009. "The Genetic Architecture of Maize Flowering Time." *Science* 325, no. 5941: 714–718. <https://doi.org/10.1126/science.1174276>.
- Burczyk, J., W. T. Adams, G. F. Moran, and A. R. Griffin. 2002. "Complex Patterns of Mating Revealed in a *Eucalyptus regnans* Seed Orchard Using Allozyme Markers and the Neighbourhood Model." *Molecular Ecology* 11, no. 11: 2379–2391. <https://doi.org/10.1046/j.1365-294X.2002.01603.x>.
- Burdon, R. D., M. J. Carson, and C. J. A. Shelbourne. 2008. "Achievements in Forest Tree Genetic Improvement in Australia and New Zealand 10: *Pinus radiata* in New Zealand." *Australian Forestry* 71, no. 4: 263–279. <https://doi.org/10.1080/00049158.2008.10675045>.
- Burns, R. M., B. H. Honkala, and [Technical coordinators]. 1990. "Silvics of North America: Volume 2. Hardwoods." In *Agriculture Handbook* 654. Washington, DC: United States Department of Agriculture (USDA), Forest Service.
- Campbell, I. D., K. McDonald, M. D. Flannigan, and J. Kringayark. 1999. "Long-Distance Transport of Pollen Into the Arctic." *Nature* 399, no. 6731: 29–30. <https://doi.org/10.1038/19891>.
- Canadell, J. G., and M. R. Raupach. 2008. "Managing Forests for Climate Change Mitigation." *Science* 320, no. 5882: 1456–1457. <https://doi.org/10.1126/science.1155458>.
- Castède, S., J. A. Campoy, J. Q. García, et al. 2014. "Genetic Determinism of Phenological Traits Highly Affected by Climate Change in *Prunus avium*: Flowering Date Dissected Into Chilling and Heat Requirements." *New Phytologist* 202, no. 2: 703–715. <https://doi.org/10.1111/nph.12658>.
- Chaix, G., O. Monteuiis, C. Garcia, et al. 2011. "Genetic Variation in Major Phenotypic Traits Among Diverse Genetic Origins of Teak (*Tectona grandis* L.f.) Planted in Taliwas, Sabah, East Malaysia." *Annals of Forest Science* 68, no. 5: 1015–1026. <https://doi.org/10.1007/s13595-011-0109-8>.
- Chakraborty, D., A. Ciceu, D. Ballian, et al. 2024. "Assisted Tree Migration Can Preserve the European Forest Carbon Sink Under Climate Change." *Nature Climate Change* 14: 845–852. <https://doi.org/10.1038/s41558-024-02080-5>.
- Chakraborty, D., N. Móricz, E. Rasztovits, L. Dobor, and S. Schueler. 2021. "Provisioning Forest and Conservation Science With High-Resolution Maps of Potential Distribution of Major European Tree Species Under Climate Change." *Annals of Forest Science* 78, no. 2: 26. <https://doi.org/10.1007/s13595-021-01029-4>.
- Clark, J. S., C. Fastie, G. Hurtt, et al. 1998. "Reid's Paradox of Rapid Plant Migration: Dispersal Theory and Interpretation of Paleoecological Records." *Bioscience* 48, no. 1: 13–24.
- Clark, P. W., A. W. D'Amato, B. J. Palik, et al. 2023. "A Lack of Ecological Diversity in Forest Nurseries Limits the Achievement of Tree-Planting Objectives in Response to Global Change." *Bioscience* 73, no. 8: 575–586. <https://doi.org/10.1093/biosci/biad049>.
- Clark, P. W., A. J. Freeman, A. W. D'Amato, et al. 2022. "Restoring a Keystone Tree Species for the Future: American Chestnut Assisted Migration Plantings in an Adaptive Silviculture Experiment." *Forest Ecology and Management* 523: 120505. <https://doi.org/10.1016/j.foreco.2022.120505>.
- Cooper, H. F., K. C. Grady, J. A. Cowan, R. J. Best, G. J. Allan, and T. G. Whitham. 2019. "Genotypic Variation in Phenological Plasticity: Reciprocal Common Gardens Reveal Adaptive Responses to Warmer Springs but Not to Fall Frost." *Global Change Biology* 25, no. 1: 187–200. <https://doi.org/10.1111/gcb.14494>.
- Critchfield, W. B. 1967. "Crossability and Relationships of the Closed-Cone Pines." *Silvae Genetica* 16: 89–97. https://www.sauerlaender-verlag.com/CMS/fileadmin/content/dokument/archiv/silvaeogenetica/16_1967/16-3-89.pdf.
- Darwin, C. 1876. *The Effects of Cross- and Self-Fertilization in the Vegetable Kingdom*. London: John Murray.
- Dauphin, B., C. Rellstab, R. O. Wüest, et al. 2023. "Re-Thinking the Environment in Landscape Genomics." *Trends in Ecology & Evolution* 38, no. 3: 261–274. <https://doi.org/10.1016/j.tree.2022.10.010>.
- Davenport, C. B. 1908. "Degeneration, Albinism and Inbreeding." *Science* 28, no. 718: 454–455. <http://www.jstor.org/stable/1634614>.
- De La Torre, A. R., D. Puiu, M. W. Crepeau, et al. 2019. "Genomic Architecture of Complex Traits in Loblolly Pine." *New Phytologist* 221, no. 4: 1789–1801. <https://doi.org/10.1111/nph.15535>.
- De la Torre, A. R., B. Wilhite, D. Puiu, et al. 2021. "Dissecting the Polygenic Basis of Cold Adaptation Using Genome-Wide Association of Traits and Environmental Data in Douglas-Fir." *Genes* 12, no. 1: 110. <https://doi.org/10.3390/genes12010110>.
- Depardieu, C., S. Gérardi, S. Nadeau, et al. 2021. "Connecting Tree-Ring Phenotypes, Genetic Associations and Transcriptomics to Decipher the Genomic Architecture of Drought Adaptation in a Widespread Conifer." *Molecular Ecology* 30, no. 16: 3898–3917. <https://doi.org/10.1111/mec.15846>.
- Dickson, R. L., G. B. Sweet, and N. D. Mitchell. 2000. "Predicting *Pinus radiata* Female Strobilus Production for Seed Orchard Site Selection in New Zealand." *Forest Ecology and Management* 133, no. 3: 197–215. [https://doi.org/10.1016/S0378-1127\(99\)00233-9](https://doi.org/10.1016/S0378-1127(99)00233-9).
- Dirlewanger, E., J. Quero-García, L. Le Dantec, et al. 2012. "Comparison of the Genetic Determinism of Two Key Phenological Traits, Flowering and Maturity Dates, in Three *Prunus* Species: Peach, Apricot and Sweet

- Cherry." *Heredity* 109, no. 5: 280–292. <https://doi.org/10.1038/hdy.2012.38>.
- Doerksen, T. K., J. Bousquet, and J. Beaulieu. 2014. "Inbreeding Depression in Intra-Provenance Crosses Driven by Founder Relatedness in White Spruce." *Tree Genetics & Genomes* 10, no. 1: 203–212. <https://doi.org/10.1007/s11295-013-0676-y>.
- Dungey, H. S. 2001. "Pine Hybrids—A Review of Their Use Performance and Genetics." *Forest Ecology and Management* 148, no. 1: 243–258. [https://doi.org/10.1016/S0378-1127\(00\)00539-9](https://doi.org/10.1016/S0378-1127(00)00539-9).
- East, E. M. 1908. "Inbreeding in Corn Reports of the Connecticut Agricultural Experimental Station for 1907." pp. 419–428.
- Eriksson, G., and B. Ilstedt. 1986. "Stem Volume of Intra- and Interprovenance Families of *Picea abies* (L.) Karst." *Scandinavian Journal of Forest Research* 1, no. 1–4: 141–152. <https://doi.org/10.1080/02827588609382407>.
- Eschen, R., A. Roques, and A. Santini. 2015. "Taxonomic Dissimilarity in Patterns of Interception and Establishment of Alien Arthropods, Nematodes and Pathogens Affecting Woody Plants in Europe." *Diversity and Distributions* 21, no. 1: 36–45. <https://doi.org/10.1111/ddi.12267>.
- Faulkner, R. 1975. *Seed Orchards, Forestry Commission Bulletin No. 54*. London: Forestry Commission Bulletin.
- Feng, L., J. Sun, Y. Shi, G. Wang, and T. Wang. 2020. "Predicting Suitable Habitats of *Camptotheca acuminata* Considering Both Climatic and Soil Variables." *Forests* 11, no. 8: 891. <https://doi.org/10.3390/f11080891>.
- Findlater, K., G. Peterson St-Laurent, S. Hagerman, and R. Kozak. 2020. "Surprisingly Malleable Public Preferences for Climate Adaptation in Forests." *Environmental Research Letters* 15: 034045. <https://doi.org/10.1088/1748-9326/ab7464>.
- Franić, I., S. Prospero, M. Hartmann, et al. 2019. "Are Traded Forest Tree Seeds a Potential Source of Nonnative Pests?" *Ecological Applications* 29, no. 7: e01971. <https://doi.org/10.1002/eap.1971>.
- Franks, S. J., and A. A. Hoffmann. 2012. "Genetics of Climate Change Adaptation." *Annual Review of Genetics* 46: 185–208. <https://doi.org/10.1146/annurev-genet-110711-155511>.
- Funda, T., and Y. A. El-Kassaby. 2012. "Seed Orchard Genetics." *CABI Reviews* 7: 1–23. <https://doi.org/10.1079/PAVSNNR2012701>.
- Gailing, O., H. Wachter, J. Heyder, H.-P. Schmitt, and R. Finkeldey. 2012. "Chloroplast DNA Analysis in Oak Stands (*Quercus robur* L.) in North Rhine-Westphalia With Presumably Slavonian Origin: Is There an Association Between Geographic Origin and Bud Phenology?" *Journal of Applied Botany and Food Quality* 81, no. 2: 165–171. <https://ojs.openagrar.de/index.php/JABFQ/issue/view/825>.
- Gapare, W. J. 2000. "Predicted and Realized Genetic Gain in *Eucalyptus grandis* Breeding Seedling Orchard in Zimbabwe." *Southern African Forestry Journal* 189, no. 1: 11–15. <https://doi.org/10.1080/10295925.2000.9631275>.
- Gorte, R. W., and P. A. Sheikh. 2010. *Deforestation and Climate Change*. Washington DC: Congressional Research Service.
- Goto, S., H. Iijima, H. Ogawa, and K. Ohya. 2011. "Outbreeding Depression Caused by Intraspecific Hybridization Between Local and Nonlocal Genotypes in *Abies sachalinensis*." *Restoration Ecology* 19, no. 2: 243–250. <https://doi.org/10.1111/j.1526-100X.2009.00568.x>.
- Grummer, J. A., T. R. Booker, R. Matthey-Doret, P. Nietlisbach, A. T. Thomaz, and M. C. Whitlock. 2022. "The Immediate Costs and Long-Term Benefits of Assisted Gene Flow in Large Populations." *Conservation Biology* 36, no. 4: e13911. <https://doi.org/10.1111/cobi.13911>.
- Harfouche, A., N. Bahrman, P. Baradat, J. P. Guyon, R. J. Petit, and A. Kremer. 2000. "Provenance Hybridization in a Diallel Mating Scheme of Maritime Pine (*Pinus pinaster*). II. Heterosis." *Canadian Journal of Forest Research* 30, no. 1: 10–16. <https://doi.org/10.1139/x99-179>.
- Harfouche, A., and A. Kremer. 2000. "Provenance Hybridization in a Diallel Mating Scheme of Maritime Pine (*Pinus pinaster*). I. Means and Variance Components." *Canadian Journal of Forest Research* 30, no. 1: 1–9. <https://doi.org/10.1139/x99-178>.
- Hazarika, R., A. Bolte, D. Bednárová, et al. 2021. "Multi-Actor Perspectives on Afforestation and Reforestation Strategies in Central Europe Under Climate Change." *Annals of Forest Science* 78, no. 3: 60. <https://doi.org/10.1007/s13595-021-01044-5>.
- Hei, N., S. Hussein, and M. Laing. 2016. "Heterosis and Combining Ability Analysis of Slow Rusting Stem Rust Resistance and Yield and Related Traits in Bread Wheat." *Euphytica* 207, no. 3: 501–514. <https://doi.org/10.1007/s10681-015-1526-9>.
- Hemery, G. E., J. R. Clark, E. Aldinger, et al. 2010. "Growing Scattered Broadleaved Tree Species in Europe in a Changing Climate: A Review of Risks and Opportunities." *Forestry: An International Journal of Forest Research* 83, no. 1: 65–81. <https://doi.org/10.1093/forestry/cpp034>.
- Hereford, J. 2009. "A Quantitative Survey of Local Adaptation and Fitness Trade-Offs." *American Naturalist* 173, no. 5: 579–588. <https://doi.org/10.1086/597611>.
- Heuchel, A., D. Hall, W. Zhao, J. Gao, U. Wennstrom, and X.-R. Wang. 2022. "Genetic Diversity and Background Pollen Contamination in Norway Spruce and Scots Pine Seed Orchard Crops." *Forest Research* 2: 1–12. <https://doi.org/10.48130/FR-2022-0008>.
- Hoegh-Guldberg, O., L. Hughes, S. McIntyre, et al. 2008. "Assisted Colonization and Rapid Climate Change." *Science* 321, no. 5887: 345–346. <https://doi.org/10.1126/science.1157897>.
- Horbach, S., R. Rauschkolb, and C. Römermann. 2023. "Flowering and Leaf Phenology Are More Variable and Stronger Associated to Functional Traits in Herbaceous Compared to Tree Species." *Flora* 300: 152218. <https://doi.org/10.1016/j.flora.2023.152218>.
- IPCC. 2022. "Climate Change 2022: Mitigation of Climate Change." In *Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by P. R. Shukla, J. Skea, R. Slade, et al. Cambridge, UK and New York, NY, USA: Cambridge University Press. <https://doi.org/10.1017/9781009157926>.
- Ivetić, V., J. Devetaković, M. Nonić, D. Stanković, and M. Šijačić-Nikolić. 2016. "Genetic Diversity and Forest Reproductive Material—From Seed Source Selection to Planting." *iForest - Biogeosciences and Forestry* 9, no. 5: 801–812. <https://doi.org/10.3832/ifor1577-009>.
- Jansen, S., and T. Geburek. 2016. "Historic Translocations of European Larch (*Larix decidua* Mill.) Genetic Resources Across Europe—A Review From the 17th Until the Mid-20th Century." *Forest Ecology and Management* 379: 114–123. <https://doi.org/10.1016/j.FORECO.2016.08.007>.
- Jansen, S., H. Konrad, and T. Geburek. 2017. "The Extent of Historic Translocation of Norway Spruce Forest Reproductive Material in Europe." *Annals of Forest Science* 74, no. 3: 56. <https://doi.org/10.1007/s13595-017-0644-z>.
- Jayawickrama, K. J. S., and L. C. Balocchi. 1993. "Tree Improvement in Chile: Two Decades of Progress." *Journal of Forestry* 91, no. 6: 43–47. <https://doi.org/10.1093/jof/91.6.43>.
- Johnsen, Ø., O. G. Dæhlen, G. Østreng, and T. Skrøppa. 2005. "Daylength and Temperature During Seed Production Interactively Affect Adaptive Performance of *Picea abies* Progenies." *New Phytologist* 168, no. 3: 589–596. <https://doi.org/10.1111/j.1469-8137.2005.01538.x>.
- Johnsen, Ø., H. Kvaalen, I. Yakovlev, O. G. Dæhlen, C. G. Fossdal, and T. Skrøppa. 2009. "An Epigenetic Memory From Time of Embryo Development Affects Climatic Adaptation in Norway Spruce." In *The 8th International Plant Cold Hardiness Seminar (8IPCH)*, 3–9. Saskatchewan, Canada: CABI Digital Library. <https://doi.org/10.1079/9781845935139.0099>.
- Johnson, I. G., G. Robinson, and A. J. O'Hara. 1992. "Wood Production and Economic Gains From the Use of Seed Orchard Stock in Radiata

- Pine Plantations in New South Wales." *Australian Forestry* 55, no. 1–4: 126–134. <https://doi.org/10.1080/00049158.1992.10676106>.
- Jones, D. F. 1917. "Dominance of Linked Factors as a Means of Accounting for Heterosis." *Genetics* 2, no. 5: 466–479. <https://doi.org/10.1093/genetics/2.5.466>.
- Jung, T., L. Orlikowski, B. Henricot, et al. 2016. "Widespread *Phytophthora* Infestations in European Nurseries Put Forest, Semi-Natural and Horticultural Ecosystems at High Risk of *Phytophthora* Diseases." *Forest Pathology* 46, no. 2: 134–163. <https://doi.org/10.1111/efp.12239>.
- Kaya, Z., and D. Lindgren. 1992. "The Genetic Variation of Inter-Provenance Hybrids of *Picea abies* and Possible Breeding Consequences." *Scandinavian Journal of Forest Research* 7, no. 1–4: 15–26. <https://doi.org/10.1080/02827589209382694>.
- Kelly, E., and B. L. Phillips. 2016. "Targeted Gene Flow for Conservation." *Conservation Biology* 30, no. 2: 259–267. <https://doi.org/10.1111/cobi.12623>.
- Konnert, M., B. Fady, D. Gömöry, et al. 2015. "Use and Transfer of Forest Reproductive Material in Europe in the Context of Climate Change." In *European Forest Genetic Resources Programme (EUFORGEN)*, 75. Rome, Italy: Bioversity International.
- Korecký, J., and Y. A. El-Kassaby. 2016. "Pollination Dynamics Variation in a Douglas-Fir Seed Orchard as Revealed by Microsatellite Analysis." *Silva Fennica* 50, no. 4: 1682. <https://doi.org/10.14214/sf.1682>.
- Kramer, K. 1995. "Phenotypic Plasticity of the Phenology of Seven European Tree Species in Relation to Climatic Warming." *Plant, Cell & Environment* 18, no. 2: 93–104. <https://doi.org/10.1111/j.1365-3040.1995.tb00356.x>.
- Langlet, O. 1971. "Two Hundred Years Genecology." *Taxon* 20, no. 5–6: 653–721. <https://doi.org/10.2307/1218596>.
- Larsen, J. 1981. "Waldbauliche und ertragskundliche Erfahrungen mit verschiedenen Provenienzen der Weißtanne (*Abies alba* Mill.) in Dänemark." *Forstwissenschaftliches Centralblatt* 100: 275–286. <https://doi.org/10.1007/BF02640644>.
- Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. "Chilling Outweighs Photoperiod in Preventing Precocious Spring Development." *Global Change Biology* 20, no. 1: 170–182. <https://doi.org/10.1111/gcb.12360>.
- Ledig, F. T., G. E. Rehfeldt, and B. Jaquish. 2012. "Projections of Suitable Habitat Under Climate Change Scenarios: Implications for Trans-Boundary Assisted Colonization." *American Journal of Botany* 99, no. 7: 1217–1230. <https://doi.org/10.3732/ajb.1200059>.
- Lefèvre, F. 2004. "Human Impacts on Forest Genetic Resources in the Temperate Zone: An Updated Review." *Forest Ecology and Management* 197, no. 1: 257–271. <https://doi.org/10.1016/j.foreco.2004.05.017>.
- Leksono, B., S. Kurinobu, and Y. Ide. 2008. "Realized Genetic Gains Observed in Second Generation Seedling Seed Orchards of *Eucalyptus pellita* in Indonesia." *Journal of Forest Research* 13, no. 2: 110–116. <https://doi.org/10.1007/s10310-008-0061-0>.
- Levkoev, E., A. Kilpeläinen, K. Luostarinen, et al. 2017. "Differences in Growth and Wood Density in Clones and Provenance Hybrid Clones of Norway Spruce." *Canadian Journal of Forest Research* 47, no. 3: 389–399. <https://doi.org/10.1139/cjfr-2016-0285>.
- Li, B., S. McKeand, and R. Weir. 1999. "Tree Improvement and Sustainable Forestry—Impact of Two Cycles of Loblolly Pine Breeding in the USA." *Forest Genetics* 6, no. 4: 229–234.
- Liebold, A. M., E. G. Brockerhoff, L. J. Garrett, J. L. Parke, and K. O. Britton. 2012. "Live Plant Imports: The Major Pathway for Forest Insect and Pathogen Invasions of the US." *Frontiers in Ecology and the Environment* 10, no. 3: 135–143. <https://doi.org/10.1890/110198>.
- Liesebach, H., K. Liepe, and C. Bäucker. 2021. "Towards New Seed Orchard Designs in Germany—A Review." *Silvae Genetica* 70, no. 1: 84–98. <https://doi.org/10.2478/sg-2021-0007>.
- Lindner, M., M. Maroschek, S. Netherer, et al. 2010. "Climate Change Impacts, Adaptive Capacity, and Vulnerability of European Forest Ecosystems." *Forest Ecology and Management* 259, no. 4: 698–709. <https://doi.org/10.1016/J.FORECO.2009.09.023>.
- Liziniewicz, M., M. Berlin, T. Solvin, et al. 2023. "Development of a Universal Height Response Model for Transfer of Norway Spruce (*Picea abies* L. Karst) in Fennoscandia." *Forest Ecology and Management* 528: 120628.
- Lstibůrek, M., M. R. García-Gil, and A. Steffenrem. 2023. "Rolling Front Landscape Breeding." *Annals of Forest Science* 80, no. 1: 36.
- Lynch, M., and B. Walsh. 1998. *Genetics and Analysis of Quantitative Traits*. Sunderland, MA: Sinauer Associates.
- Madhibha, T., R. Murepa, C. Musokonyi, and W. Gapare. 2013. "Genetic Parameter Estimates for Interspecific *Eucalyptus* Hybrids and Implications for Hybrid Breeding Strategy." *New Forests* 44, no. 1: 63–84. <https://doi.org/10.1007/s11056-011-9302-8>.
- Marchal, A., F. Muñoz, F. Millier, L. Sánchez, and L. E. Pâques. 2017. "Hybrid Larch Heterosis: For Which Traits and Under Which Genetic Control?" *Tree Genetics & Genomes* 13, no. 5: 92. <https://doi.org/10.1007/s11295-017-1177-1>.
- McKone, M. J., and D. L. Hernández. 2021. "Community-Level Assisted Migration for Climate-Appropriate Prairie Restoration." *Restoration Ecology* 29, no. 7: e13416. <https://doi.org/10.1111/rec.13416>.
- Milesi, P., M. Berlin, J. Chen, et al. 2019. "Assessing the Potential for Assisted Gene Flow Using Past Introduction of Norway Spruce in Southern Sweden: Local Adaptation and Genetic Basis of Quantitative Traits in Trees." *Evolutionary Applications* 12, no. 10: 1946–1959. <https://doi.org/10.1111/eva.12855>.
- Miller, L. K., and J. DeBell. 2013. "Current Seed Orchard Techniques and Innovations." In *National Proceedings: Forest and Conservation Nursery Associations – 2012. Proceedings RMRS-P-69*, edited by D. L. Haase, J. R. Pinto, K. M. Wilkinson, and Technical Coordinators, 80–86. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Missanjo, E., G. Kamanga-Thole, and V. Manda. 2013. "Estimation of Genetic and Phenotypic Parameters for Growth Traits in a Clonal Seed Orchard of *Pinus Kesiya* in Malawi." *International Scholarly Research Notices* 2013, no. 1: 346982. <https://doi.org/10.1155/2013/346982>.
- Moll, R. H., J. H. Lonnquist, J. V. Fortuno, and E. C. Johnson. 1965. "The Relationship of Heterosis and Genetic Divergence in Maize." *Genetics* 52, no. 1: 139–144. <https://doi.org/10.1093/genetics/52.1.139>.
- Montwé, D., M. Isaac-Renton, A. Hamann, and H. Spiecker. 2018. "Cold Adaptation Recorded in Tree Rings Highlights Risks Associated With Climate Change and Assisted Migration." *Nature Communications* 9, no. 1: 1574. <https://doi.org/10.1038/s41467-018-04039-5>.
- Moriguchi, Y., N. Tani, S. Ito, et al. 2005. "Gene Flow and Mating System in Five *Cryptomeria japonica* D. Don Seed Orchards as Revealed by Analysis of Microsatellite Markers." *Tree Genetics & Genomes* 1, no. 4: 174–183. <https://doi.org/10.1007/s11295-005-0023-z>.
- Mueller, J. M., and J. J. Hellmann. 2008. "An Assessment of Invasion Risk From Assisted Migration." *Conservation Biology* 22, no. 3: 562–567. <https://doi.org/10.1111/j.1523-1739.2008.00952.x>.
- Müller, F. 1997. "Walderhaltung in Oesterreich: Zehn Jahre Sicherung forstlicher Genressourcen." *LÖBF-Mitteilungen* 22, no. 4: 50–53.
- Myking, T., M. Rusanen, A. Steffenrem, E. D. Kjær, and G. Jansson. 2016. "Historic Transfer of Forest Reproductive Material in the Nordic Region: Drivers, Scale and Implications." *Forestry: An International Journal of Forest Research* 89, no. 4: 325–337. <https://doi.org/10.1093/forestry/cpw020>.
- Na, S.-J., H.-S. Lee, S.-U. Han, J.-M. Park, and K.-S. Kang. 2015. "Estimation of Genetic Gain and Diversity Under Various Genetic Thinning Scenarios in a Breeding Seed Orchard of *Quercus acutissima*."

- Scandinavian Journal of Forest Research 30, no. 5: 377–381. <https://doi.org/10.1080/02827581.2015.1018936>.
- Nathan, R., N. Horvitz, Y. He, A. Kuparinen, F. M. Schurr, and G. G. Katul. 2011. "Spread of North American Wind-Dispersed Trees in Future Environments." *Ecology Letters* 14, no. 3: 211–219. <https://doi.org/10.1111/j.1461-0248.2010.01573.x>.
- Nathan, R., G. G. Katul, H. S. Horn, et al. 2002. "Mechanisms of Long-Distance Dispersal of Seeds by Wind." *Nature* 418, no. 6896: 409–413. <https://doi.org/10.1038/nature00844>.
- Neophytou, C., M. van Loo, and H. Hasenauer. 2020. "Genetic Diversity in Introduced Douglas-Fir and Its Natural Regeneration in Central Europe." *Forestry: An International Journal of Forest Research* 93, no. 4: 535–544. <https://doi.org/10.1093/forestry/cpz055>.
- Ni, M., and M. Vellend. 2024. "Soil Properties Constrain Predicted Poleward Migration of Plants Under Climate Change." *New Phytologist* 241, no. 1: 131–141. <https://doi.org/10.1111/nph.19164>.
- Nicodemus, A., M. Varghese, B. Nagarajan, and D. Lindgren. 2009. "Annual Fertility Variation in Clonal Seed Orchards of Teak (*Tectona grandis* L.F.) and Its Impact on Seed Crop." *Silvae Genetica* 58, no. 1–6: 85–93. <https://doi.org/10.1515/sg-2009-0011>.
- Nielsen, U. B., J. Xu, and O. K. Hansen. 2020. "Genetics in and Opportunities for Improvement of Nordmann Fir (*Abies nordmanniana* (Steven) Spach) Christmas Tree Production." *Tree Genetics & Genomes* 16, no. 5: 66. <https://doi.org/10.1007/s11295-020-01461-z>.
- O'Hara, K. L. 2016. "What Is Close-To-Nature Silviculture in a Changing World?" *Forestry: An International Journal of Forest Research* 89, no. 1: 1–6. <https://doi.org/10.1093/forestry/cpv043>.
- Pearse, I. S., W. D. Koenig, K. A. Funk, and M. B. Pesendorfer. 2015. "Pollen Limitation and Flower Abortion in a Wind-Pollinated, Masting Tree." *Ecology* 96, no. 2: 587–593. <https://doi.org/10.1890/14-0297.1>.
- Petit, R. J., I. Aguinalalde, J.-L. de Beaulieu, et al. 2003. "Glacial Refugia: Hotspots but Not Melting Pots of Genetic Diversity." *Science* 300, no. 5625: 1563–1565. <https://doi.org/10.1126/science.1083264>.
- Powers, L. 1944. "An Expansion of Jones's Theory for the Explanation of Heterosis." *American Naturalist* 78, no. 776: 275–280. <https://doi.org/10.1086/281199>.
- Rehfeldt, G. E., N. M. Tchebakova, Y. I. Parfenova, W. R. Wyckoff, N. A. Kuzmina, and L. I. Milyutin. 2002. "Intraspecific Responses to Climate in *Pinus sylvestris*." *Global Change Biology* 8, no. 9: 912–929. <https://doi.org/10.1046/j.1365-2486.2002.00516.x>.
- Reid, C. 1899. *The Origin of the British Flora*. London: Dulau.
- Reid, D. J. S. 2008. "British Columbia's Seed Orchard Program: Multi Species Management With Integration to the End User." In *Proceedings of a Seed Orchard Conference, Umeå, Sweden, 26–28 September 2007*, edited by D. Lindgren, 204–214. Sweden: Swedish University of Agricultural Sciences.
- Richardson, D. M., J. J. Hellmann, J. S. McLachlan, et al. 2009. "Multidimensional Evaluation of Managed Relocation." *Proceedings of the National Academy of Sciences* 106, no. 24: 9721–9724. <https://doi.org/10.1073/pnas.0902327106>.
- Riehl, J. F. L., C. T. Cole, C. J. Morrow, et al. 2023. "Genomic and Transcriptomic Analyses Reveal Polygenic Architecture for Ecologically Important Traits in Aspen (*Populus tremuloides* Michx.)." *Ecology and Evolution* 13, no. 10: e10541. <https://doi.org/10.1002/ece3.10541>.
- Rockwood, D. L., D. A. Huber, and T. L. White. 2001. "Provenance and Family Variability in Slash Pine (*Pinus elliottii* var. *elliottii* Engelm.) Grown in Southern Brazil and Northeastern Argentina." *New Forests* 21, no. 2: 115–125. <https://doi.org/10.1023/A:1011850009675>.
- Sáenz-Romero, C., R. A. Lindig-Cisneros, D. G. Joyce, J. Beaulieu, J. B. St Clair, and B. C. Jaquish. 2016. "Assisted Migration of Forest Populations for Adapting Trees to Climate Change." *Revista Chapingo Serie Ciencias Forestales y Del Ambiente* 22, no. 3: 303–323. <https://doi.org/10.5154/rchscfa.2014.10.052>.
- Sáenz-Romero, C., G. Neill, S. N. Aitken, and R. Lindig-Cisneros. 2021. "Assisted Migration Field Tests in Canada and Mexico: Lessons, Limitations, and Challenges." *Forests* 12, no. 1: 9. <https://doi.org/10.3390/f12010009>.
- Saha, S., C. Kuehne, U. Kohnle, et al. 2012. "Growth and Quality of Young Oaks (*Quercus Robur* and *Quercus petraea*) Grown in Cluster Plantings in Central Europe: A Weighted Meta-Analysis." *Forest Ecology and Management* 283: 106–118. <https://doi.org/10.1016/j.foreco.2012.07.021>.
- Sánchez-Pérez, R., J. Del Cueto, F. Dicenta, and P. Martínez-Gómez. 2014. "Recent Advancements to Study Flowering Time in Almond and Other *Prunus* Species." *Frontiers in Plant Science* 5: 334. <https://doi.org/10.3389/fpls.2014.00334>.
- Savolainen, O., T. Pyhäjärvi, and T. Knürr. 2007. "Gene Flow and Local Adaptation in Trees." *Annual Review of Ecology, Evolution, and Systematics* 38: 595–619. <https://doi.org/10.1146/annurev.ecolsys.38.091206.095646>.
- Schreiber, S. G., U. G. Hacke, and A. Hamann. 2015. "Variation of Xylem Vessel Diameters Across a Climate Gradient: Insight From a Reciprocal Transplant Experiment With a Widespread Boreal Tree." *Functional Ecology* 29, no. 11: 1392–1401. <https://doi.org/10.1111/1365-2435.12455>.
- Secretariat of the Convention on Biological Diversity. 2011. "Nagoya Protocol on Access to Genetic Resources and the Fair and Equitable Sharing of Benefits Arising From Their Utilization to the Convention on Biological Diversity." <https://www.cbd.int/abs/doc/protocol/nagoya-protocol-en.pdf>.
- Shelton, W. R., R. J. Mitchell, D. A. Christopher, L. P. Jack, and J. D. Karron. 2024. "Among-Individual Variation in Flowering Phenology Affects Flowering Synchrony and Mating Opportunity." *American Journal of Botany* 111, no. 1: e16269. <https://doi.org/10.1002/ajb2.16269>.
- Shi, T., R. J. Arnold, W. Kang, et al. 2016. "Genetic Variation and Gains for Two Generations of *Eucalyptus dunnii* in China." *Australian Forestry* 79, no. 1: 15–24. <https://doi.org/10.1080/00049158.2015.1086720>.
- Shull, G. H. 1908. "The Composition of a Field of Maize." *Journal of Heredity* 1: 296–301.
- Shull, G. H. 1914. "Duplicate Genes for Capsule-Form in *Bursa bursa-pastoris*." *Zeitschrift für Induktive Abstammungs- und Vererbungslehre* 12, no. 1: 97–149. <https://doi.org/10.1007/BF01837282>.
- Silvestro, R., S. Rossi, S. Zhang, I. Froment, J. G. Huang, and A. Saracino. 2019. "From Phenology to Forest Management: Ecotypes Selection Can Avoid Early or Late Frosts, but Not Both." *Forest Ecology and Management* 436: 21–26. <https://doi.org/10.1016/j.foreco.2019.01.005>.
- Singh, P., J. B. St Clair, B. M. Lind, et al. 2024. "Genetic Architecture of Disease Resistance and Tolerance in Douglas-Fir Trees." *New Phytologist* 243, no. 2: 705–719. <https://doi.org/10.1111/nph.19797>.
- Sittaro, F., A. Paquette, C. Messier, and C. A. Nock. 2017. "Tree Range Expansion in Eastern North America Fails to Keep Pace With Climate Warming at Northern Range Limits." *Global Change Biology* 23, no. 8: 3292–3301. <https://doi.org/10.1111/gcb.13622>.
- Skogforsk. 2024. "Planter's Guide." <https://www.skogforsk.se/plantersguide>.
- Smith, C., J. C. A. Baker, and D. V. Spracklen. 2023. "Tropical Deforestation Causes Large Reductions in Observed Precipitation." *Nature* 615, no. 7951: 270–275. <https://doi.org/10.1038/s41586-022-05690-1>.
- So, T., I. Theilade, and B. Dell. 2010. "Conservation and Utilization of Threatened Hardwood Species Through Reforestation? An Example of *Azelia Xylocarpa* (Kruz.) Craib and *Dalbergia cochinchinensis* Pierre in Cambodia." *Pacific Conservation Biology* 16, no. 2: 101–116. <https://www.publish.csiro.au/PC/PC100101>.

- Song, J., B. Ratcliffe, T. Kess, B. S. Lai, J. Korecký, and Y. A. El-Kassaby. 2018. "Temporal Quantification of Mating System Parameters in a Coastal Douglas-Fir Seed Orchard Under Manipulated Pollination Environment." *Scientific Reports* 8, no. 1: 11593. <https://doi.org/10.1038/s41598-018-30041-4>.
- Stejskal, J., J. Horák, and J. Typta. 2016. "Effect of Hybridization in the Firs: Artificial Hybridization May Lead to Higher Survival Rate." *European Journal of Forest Research* 135: 1097–1105. <https://doi.org/10.1007/s10342-016-0996-1>.
- Ste-Marie, C., A. Nelson, A. Dabros, and M.-E. Bonneau. 2011. "Assisted Migration: Introduction to a Multifaceted Concept." *Forestry Chronicle* 87, no. 6: 724–730. <https://doi.org/10.5558/tfc2011-089>.
- Stinziano, J. R., and D. A. Way. 2014. "Combined Effects of Rising [CO₂] and Temperature on Boreal Forests: Growth, Physiology and Limitations." *Botany* 92, no. 6: 425–436. <https://doi.org/10.1139/cjb-2013-0314>.
- Swain, T.-L., S. D. Verry, and M. D. Laing. 2013. "A Comparison of the Effect of Genetic Improvement, Seed Source and Seedling Seed Orchard Variables on Progeny Growth in *Eucalyptus nitens* in South Africa." *Tree Genetics & Genomes* 9, no. 3: 767–778. <https://doi.org/10.1007/s11295-013-0593-0>.
- Tabel, U. 1997. "Erhaltung forstlicher Genressourcen in der Bundesrepublik Deutschland." *AFZ/Der Wald* 52, no. 5: 259–261.
- Turner, J. A. 1997. "Realised Genetic Gain in *Pinus radiata* From "850" Seed-Orchard Seedlots Grown Commercially in the Central North Island, New Zealand. Part 1: Growth." *New Zealand Journal of Forestry Science* 27, no. 2: 142–157.
- Urrestarazu, J., H. Muranty, C. Denancé, et al. 2017. "Genome-Wide Association Mapping of Flowering and Ripening Periods in Apple." *Frontiers in Plant Science* 8: 1923. <https://doi.org/10.3389/fpls.2017.01923>.
- Van Daele, F., O. Honnay, and H. De Kort. 2022. "Genomic Analyses Point to a Low Evolutionary Potential of Prospective Source Populations for Assisted Migration in a Forest Herb." *Evolutionary Applications* 15, no. 11: 1859–1874. <https://doi.org/10.1111/eva.13485>.
- Vettraino, A. M., R. Potting, and R. Raposo. 2018. "EU Legislation on Forest Plant Health: An Overview With a Focus on *Fusarium circinatum*." *Forests* 9, no. 9: 568. <https://doi.org/10.3390/f9090568>.
- Visser, M. E. 2008. "Keeping Up With a Warming World; Assessing the Rate of Adaptation to Climate Change." *Proceedings of the Royal Society B: Biological Sciences* 275, no. 1635: 649–659.
- Wadgyamar, S. M., M. N. Cumming, and A. E. Weis. 2015. "The Success of Assisted Colonization and Assisted Gene Flow Depends on Phenology." *Global Change Biology* 21, no. 10: 3786–3799. <https://doi.org/10.1111/gcb.12988>.
- Wadgyamar, S. M., and A. E. Weis. 2017. "Phenological Mismatch and the Effectiveness of Assisted Gene Flow." *Conservation Biology* 31, no. 3: 547–558. <https://doi.org/10.1111/cobi.12877>.
- Way, D. A., and R. A. Montgomery. 2015. "Photoperiod Constraints on Tree Phenology, Performance and Migration in a Warming World." *Plant, Cell & Environment* 38, no. 9: 1725–1736. <https://doi.org/10.1111/pce.12431>.
- Weis, A. E. 2015. "On the Potential Strength and Consequences for Nonrandom Gene Flow Caused by Local Adaptation in Flowering Time." *Journal of Evolutionary Biology* 28, no. 3: 699–714. <https://doi.org/10.1111/jeb.12612>.
- West-Eberhard, M. J. 1989. "Phenotypic Plasticity and the Origins of Diversity." *Annual Review of Ecology and Systematics* 20, no. 1: 249–278. <https://doi.org/10.1146/annurev.es.20.110189.001341>.
- White, T. L., W. T. Adams, and D. B. Neale. 2007. *Forest Genetics*. Cambridge MA: CABI Publishing.
- Whitlock, M. C., P. K. Ingvarsson, and T. Hatfield. 2000. "Local Drift Load and the Heterosis of Interconnected Populations." *Heredity* 84, no. 4: 452–457. <https://doi.org/10.1046/j.1365-2540.2000.00693.x>.
- Whitlock, R., G. B. Stewart, S. J. Goodman, et al. 2013. "A Systematic Review of Phenotypic Responses to Between-Population Outbreeding." *Environmental Evidence* 2, no. 1: 13. <https://doi.org/10.1186/2047-2382-2-13>.
- Wu, H. X., K. G. Eldridge, A. C. Matheson, et al. 2007. "Achievements in Forest Tree Improvement in Australia and New Zealand 8. Successful Introduction and Breeding of Radiata Pine in Australia." *Australian Forestry* 70, no. 4: 215–225. <https://doi.org/10.1080/00049158.2007.10675023>.
- Würschum, T., X. Zhu, Y. Zhao, Y. Jiang, J. C. Reif, and H. P. Maurer. 2023. "Maximization Through Optimization? On the Relationship Between Hybrid Performance and Parental Genetic Distance." *Theoretical and Applied Genetics* 136, no. 9: 186. <https://doi.org/10.1007/s00122-023-04436-5>.
- Xu, J., U. B. Nielsen, and O. K. Hansen. 2018. "Ad Hoc Breeding of *Abies bornmülleriana* for Christmas Tree Production Using a Combination of DNA Markers and Quantitative Genetics—A Case Study." *Tree Genetics & Genomes* 14, no. 5: 64. <https://doi.org/10.1007/s11295-018-1276-7>.
- Xu, W., and C. E. Prescott. 2024. "Can Assisted Migration Mitigate Climate-Change Impacts on Forests?" *Forest Ecology and Management* 556: 121738. <https://doi.org/10.1016/j.foreco.2024.121738>.
- Xu, W., J. M. Rhemtulla, D. Luo, and T. Wang. 2024. "Common Drivers Shaping Niche Distribution and Climate Change Responses of One Hundred Tree Species." *Journal of Environmental Management* 370: 123074. <https://doi.org/10.1016/j.jenvman.2024.123074>.
- Yu, D., X. Gu, S. Zhang, et al. 2021. "Molecular Basis of Heterosis and Related Breeding Strategies Reveal Its Importance in Vegetable Breeding." *Horticulture Research* 8: 120. <https://doi.org/10.1038/s41438-021-00552-9>.
- Zanewich, K. P., D. W. Pearce, and S. B. Rood. 2018. "Heterosis in Poplar Involves Phenotypic Stability: Cottonwood Hybrids Outperform Their Parental Species at Suboptimal Temperatures." *Tree Physiology* 38, no. 6: 789–800. <https://doi.org/10.1093/treephys/tpy019>.
- Zhu, K., C. W. Woodall, and J. S. Clark. 2012. "Failure to Migrate: Lack of Tree Range Expansion in Response to Climate Change." *Global Change Biology* 18, no. 3: 1042–1052. <https://doi.org/10.1111/j.1365-2486.2011.02571.x>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.